

[Get Access](#)[Share](#)[Export](#)

## Immunology Letters

Volume 197, May 2018, Pages 1-8

# Human IgG subclass cross-species reactivity to mouse and cynomolgus monkey Fcγ receptors

Mehabaw G. Derebe  , Rupesh K. Nanjunda, Gary L. Gilliland, Eilyn R. Lacy, Mark L. Chiu

 [Show more](#)

<https://doi.org/10.1016/j.imlet.2018.02.006>

[Get rights and content](#)

### Highlights

- Affinities of human IgG subclasses to mouse and cynomolgus monkey Fcγ receptors presented
- Cyno FcγRI had nanomolar affinity to human IgG4 PAA while human FcγRI bound weakly.
- Cyno FcγRI had nanomolar affinity to human IgG4 PAA while human FcγRI bound weakly.

**Register to receive personalized recommendations based on your recent signed-in activity** 

Register for free

### Abstract

In therapeutic antibody discovery and early development, mice and cynomolgus monkey are used as animal models to assess toxicity, efficacy and other properties of candidate molecules. As more candidate antibodies are based on human immunoglobulin (IgG) subclasses, many strategies are pursued to simulate the human system in the test animal. However, translation rate from a successful preclinical trial to an approved drug is extremely low. This may partly be due to differences in interaction of human IgG based candidate molecules to endogenous Fcγ receptors of model animals in comparison to those of human Fcγ receptors. In this study, we compare binding characteristics of human IgG subclasses commonly used in drug development (IgG1, IgG2, IgG4) and their respective Fc silent versions (IgG1σ, IgG2σ, IgG4 PAA) to human, mouse, and cynomolgus monkey Fcγ

receptors. To control interactions between Fab and Fc domains, the test IgGs all have the same variable region sequences. We found distinct variations of interaction of human IgG subclasses to model animal Fcγ receptors in comparison to their human counterparts. Particularly, cynomolgus monkey Fcγ receptors showed consistently tighter binding to human IgGs than human Fcγ receptors. Moreover, the presumably Fc silent human IgG4 PAA framework bound to cynomolgus monkey FcγRI with nanomolar affinity while only very weak binding was observed for the human FcγRI. Our results highlighted the need for a thorough *in vitro* affinity characterization of candidate IgGs against model animal Fcγ receptors and careful design of preclinical studies.

 [Previous](#)

[Next](#) 

## Keywords

Immunoglobulin G (IgG); Fcγ receptors; Animal models; Cross-species reactivity; Fc-silent IgGs; Surface Plasmon Resonance (SPR)

---

[Recommended articles](#)

[Citing articles \(1\)](#)

[View full text](#)

© 2018 Published by Elsevier B.V. on behalf of European Federation of Immunological Societies.

---

**ELSEVIER**

[About ScienceDirect](#) [Remote access](#) [Shopping cart](#) [Contact and support](#)  
[Terms and conditions](#) [Privacy policy](#)

We use cookies to help provide and enhance our service and tailor content and ads. By continuing you agree to the [use of cookies](#).

Copyright © 2018 Elsevier B.V. or its licensors or contributors. ScienceDirect® is a registered trademark of Elsevier B.V.

 RELX Group™